



A case of bilateral perinatal testicular torsion that presented with unilateral torsion; necessity of contralateral testis exploration

Hamid Pakmanesh , Mohsen Alinejad 

Cite this article as: Pakmanesh H, Alinejad M. A case of bilateral perinatal testicular torsion that presented with unilateral torsion; necessity of contralateral testis exploration. Turk J Urol 2018; 44(6): 511-4.

ABSTRACT

Perinatal testicular torsion is a rare emergency in a neonate that prompts immediate attention. Bilateral testicular torsion is extremely rare. We report a case of bilateral torsion that presented with unilateral scrotal swelling but significant atrophy and dark discoloration of the contralateral testis that was secondary to asynchronous prenatal torsion. There is no consensus about exploration of the contralateral testis when exploring a case with unilateral testicular torsion. Nevertheless, findings in this case report indicate that bilateral exploration is mandatory in each case of perinatal testicular torsion to evaluate the condition of contralateral testis and fix it to prevent development of future torsion that may result in anorchia.

Keywords: Neonatal; testicular; torsion.

Introduction

Perinatal testicular torsion is defined as testicular torsion that is diagnosed at birth (prenatal testicular torsion) or occurs during the first month of life (neonatal testicular torsion).^[1,2] This is a rare entity and its etiology is not well understood, however some associations have been suggested, including preeclampsia, gestational diabetes, twin gestation, large size for gestational age, presence of prenatal hydronephrosis or prolonged delivery.^[2,3] In addition, there may be a role for genetic factors, as this entity has been also reported in siblings.^[4,5]

It has been suggested that exclusively extravaginal twisting of the spermatic cord which is due to poor adhesion of the tunica vaginalis to the gubernaculum involves in the pathogenesis of the perinatal torsion. As a result, testis and tunica vaginalis twist together as a unit and induces testicular asphyxia.^[3,6,7] Despite that, intravaginal testicular torsion is associated with a typical deformity in which testis is not fixed in

the tunica vaginalis and can twist freely on its pedicle like a a bell clapper. This type of torsion is called intravaginal torsion which is frequently seen in prepubertal boys. Intravaginal torsion has been reported in an undescended testis of a 30-day newborn boy.^[8]

Although previously emergent exploration in all cases with neonatal torsion was recommended,^[1,9,10] the advantage of this approach for the newborn baby has been questioned recently.^[2] Nevertheless, we report a case with unilateral presentation of testicular swelling and pain with normal appearing contralateral testis that was finally diagnosed as bilateral torsion during exploration. This case supports emergent exploration in these cases to evaluate the condition of contralateral testis and fix it to prevent future torsion that may end in anorchia.

Case presentation

A boy was referred to our emergency department for urologic consult at postnatal 12th hour. He

ORCID IDs of the authors:

H.P. 0000-0001-5534-7042;
M.A. 0000-0002-8282-4892

Department of Urology, Shahid Bahonar Hospital, Kerman University of Medical Sciences (KMU), Kerman, Iran

Submitted:
13.10.2016

Accepted:
23.08.2017

Available Online Date:
28.08.2018

Corresponding Author:
Hamid Pakmanesh
E-mail:
h_pakmanesh@kmu.ac.ir

©Copyright 2018 by Turkish Association of Urology

Available online at
www.turkishjournalofurology.com

Table 1. Results of some reports of perinatal testicular torsion

Author, date	Number of cases	Presentation	Result of exploration	Decision for symptomatic side	Contralateral testis	Follow up	Recommendation for emergency exploration	Recommendation for contralateral testis exploration
Van Glabeke et al. ^[16] , 2000	18	Unilateral	Unilateral	Orchidectomy in 17 cases, one case testes preserved	Pexed		Yes	Yes
Olguner et al. ^[17] , 2000	1	Bilateral	Bilateral	Right testis Orchidectomy and left testis detorsion			Yes	Yes
Arena et al. ^[18] , 2005	1	Bilateral	Bilateral	Two testis detorsion			Yes	Yes
Al-Salem AH. ^[19] , 2007	11	Unilateral	10 cases Unilateral and 1 bilateral	In 7 cases orchidectomy and other testes preserved and follow	Pexed	Ischemic	Yes Preserved testes	Yes
Ahmed et al. ^[20] , 2008	2	Unilateral	Bilateral*		Pexed		Yes	Yes
Ralahy et al. ^[21] , 2010	1	Unilateral	Unilateral	Orchidectomy	Pexed		Yes	Yes
Sachwitz et al. ^[22] , 2012	1	Bilateral	Bilateral	Bilateral orchidectomy	Pexed		Yes	Yes
Drlik et al. ^[13] , 2013	1	Unilateral	Bilateral*	Unilateral orchidectomy	Pexed	Necrosis of the	Yes Preserved testis	Yes
Granger et al. ^[14] , 2016	2	Unilateral	Bilateral*	Unilateral orchidectomy in two cases	Pexed	No atrophy in	Preserved testis	Yes
Abraham et al. ^[12] , 2016	28	Unilateral	3 bilateral cases*			Pexed		Yes
Kawamura et al. ^[23] , 2016	1	Unilateral	Unilateral	Orchidectomy	Pexed		Yes	Yes

*Incidental finding during exploration of a case with contralateral testis torsion

weighed 3.600-g, and, the second baby of a 30-year-old woman (gravida 2, para 2) with history of gestational DM and hypothyroidism. Routine prenatal sonogram obtained at the 36th week of gestation was normal, so repeat cesarean section had been planned and performed at the 38th week+2days of gestation.

Pediatrician referred the neonate to us because of erythema and swelling of the left side of the scrotum associated with agitation and persistent crying. In physical examination, the scrotal wall was erythematous at the left side, the left testis was tender and bigger than normal, and we were not able to differentiate the epididymal margin. The right testis was small but normal in physical examination. Sonographic evaluation showed heterogeneous pattern of the left testis and mild hydrocele at the right side. Further, both testes showed no perfusion in color Doppler sonography. The patient underwent surgical exploration immediately. The left testis was edematous and necrotic and a 1080° extravaginal torsion was apparent (Figure 1). The right testis was atrophic and dark colored because of recurrent prenatal intravaginal torsions. The testis had no connection to the scrotal soft tissue that is the typical bell clapper deformity (Figure 2). The left necrotic testis was removed. The right atrophic testis was preserved because of possible endocrine function and was fixed to scrotal wall to prevent torsion. The postoperative course was uneventful. After four months, the right testis was completely normal in physical examination. Written informed consent was obtained from parents of the patient for publication of this case report and relevant images.



Figure 1. The left testis was edematous and necrotic and a 1080° extravaginal torsion was apparent



Figure 2. The right testis was atrophic and dark colored because of recurrent prenatal torsion

Discussion

In case of perinatal testicular torsion, surgeon should weigh the risk of anesthesia in a newborn against the possibility of pre-

serving testicular function. Previously most specialists validated early surgical intervention,^[1,9,10] while recently some experts do not recommend emergent exploration for these patients since in their experience none of the cases had derived any benefit from early exploration.^[2] Nevertheless, boys with severe pain and agitation associated with scrotal swelling and erythema and tenderness favouring recent testicular torsion, like our patient, should undergo testicular exploration to relieve pain, to assess testis viability and to fix contralateral testis so as to prevent subsequent anorchia.

Regarding exploration of the contralateral testis, in a survey performed in the United Kingdom and Ireland, 21 percent of the surgeons responded that they had not performed contralateral orchidopexy with concerns of damaging a healthy testis.^[11] Conversely, Ahmed et al presented two cases with swelling of unilateral testis in newborns with normally appearing contralateral testis in whom bilateral testicular torsion was observed during surgical exploration. To the best of our knowledge, three case reports including six similar patients have been reported since 2008 (Table 1).^[12-14,16-23] This finding also indicates the importance of exploration of the contralateral testis even with normal preoperative findings. This is especially important when the twisted testis has been removed.

Although sonography reported mild contralateral hydrocele, we explored the contralateral testis from a scrotal incision to prevent unwanted vasal or cord injury. Kaefer et al.^[15] reported follow-up of 37 cases of neonatal testicular torsion with contralateral hydrocele that were explored with scrotal approach, in whom possibly patent processus vaginalis was present. They did not encounter any case of hernia recurrence during 1-14 years follow-up.

Full examination of the genitalia should be performed in each neonate. Bilateral exploration is mandatory in each case of perinatal torsion to evaluate the condition of contralateral testis and fix it to prevent future torsion that may end in anorchia.

Informed Consent: Written informed consent was obtained from parents of the boy for publication of this case report and relevant images.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept – H.P.; Design – H.P.; Supervision – H.P.; Resources – H.P., M.A.; Materials – M.A.; Data Collection and/or Processing – M.A., H.P.; Analysis and/or Interpretation – H.P., M.A.; Literature Search – H.P., M.A.; Writing Manuscript – M.A., H.P.; Critical Review – H.P., M.A.

Conflict of Interest: Authors have no conflicts of interest to declare.

Financial Disclosure: The authors have declared that they didn't receive any financial support for this study.

References

1. Brandt MT, Sheldon CA, Wacksman J, Matthews P. Prenatal testicular torsion: principles of management. *J Urol* 1992;147:670-2. [\[CrossRef\]](#)
2. Kaye JD, Levitt SB, Friedman SC, Franco I, Gitlin J, Palmer LS. Neonatal torsion: a 14-year experience and proposed algorithm for management. *J Urol* 2008;179:2377-83. [\[CrossRef\]](#)
3. Riaz-Ul-Haq M, Mahdi DE, Elhassan EU. Neonatal testicular torsion; a review article. *Iran J Pediatr* 2012;22:281-9.
4. Gorbonos A, Cheng EY. Perinatal testicular torsion in siblings. *J Pediatr Urol* 2007;3:514-5. [\[CrossRef\]](#)
5. Castilla EE, Sod R, Anzorena O, Texido J. Neonatal testicular torsion in two brothers. *J Med Genet* 1975;12:112-3. [\[CrossRef\]](#)
6. Driver CP, Losty PD. Neonatal testicular torsion. *Br J Urol* 1998;82:855-8. [\[CrossRef\]](#)
7. Gillenwater JY, Burros HM. Torsion of the spermatic cord in utero. *JAMA* 1966;198:1123-4. [\[CrossRef\]](#)
8. Fernandez MS, Dominguez C, Lopez A, Benlloch C, Garcia-Ibarra F. Neonatal intravaginal testicular torsion in an undescended testis. *Cir Pediatr* 1996;9:128-9.
9. Arnbjornsson E. Testicular survival after neonatal torsion. *Z Kinderchir* 1986;41:293-4. [\[CrossRef\]](#)
10. Pinto KJ, Noe HN, Jerkins GR. Management of neonatal testicular torsion. *J Urol* 1997;158:1196-7. [\[CrossRef\]](#)
11. Rhodes HL, Corbett HJ, Horwood JF, Losty PD. Neonatal testicular torsion: a survey of current practice amongst paediatric surgeons and urologists in the United Kingdom and Ireland. *J Pediatr Surg* 2011;46:2157-60. [\[CrossRef\]](#)
12. Abraham MB, Charles A, Gera P, Srinivasjois R. Surgically managed perinatal testicular torsion: a single centre experience. *J Matern Fetal Neonatal Med* 2016;29:1265-8. [\[CrossRef\]](#)
13. Drlik M, Kocvara R. Bilateral perinatal torsion of spermatic cord - a case report and literature review. *Rozhl Chir* 2013;92:98-101.
14. Granger J, Brownlee EM, Cundy TP, Goh DW. Bilateral perinatal testicular torsion: successful salvage supports emergency surgery. *BMJ Case Rep* 2016;pii: bcr2016216020.
15. Kaefer M, Agarwal D, Misseri R, Whittam B, Hubert K, Szyman-ski K, et al. Treatment of contralateral hydrocele in neonatal testicular torsion: Is less more? *J Pediatr Urol* 2016;306.e1-306.e4.
16. Van Glabeke E, Philippe-Chomette P, Gall O, Oro H, Larroquet M, Audry G. Spermatic cord torsion in the newborn: role of surgical exploration. *Arch Pediatr* 2000;7:1072-6. [\[CrossRef\]](#)
17. Olguner M, Akgur FM, Aktug T, Derebek E. Bilateral asynchronous perinatal testicular torsion: a case report. *J Pediatr Surg* 2000;35:1348-9. [\[CrossRef\]](#)
18. Arena F, Nicotina PA, Scalfari G, Visalli C, Arena S, Zuccarello B, et al. A case of bilateral prenatal testicular torsion: Ultrasonographic features, histopathological findings and management. *J Pediatr Urol* 2005;1:369-72. [\[CrossRef\]](#)
19. Al-Salem AH. Intrauterine testicular torsion: a surgical emergency. *J Pediatr Surg* 2007;42:1887-91. [\[CrossRef\]](#)
20. Ahmed SJ, Kaplan GW, DeCambre ME. Perinatal testicular torsion: preoperative radiological findings and the argument for urgent surgical exploration. *J Pediatr Surg* 2008;43:1563-5. [\[CrossRef\]](#)
21. Ralahy FM, Rambel H, Rakototiana FA, Rabenasolo M, Rantomalala Y, Andriamanarivo L. A case of extravaginal twisting of spermatic cord. *Arch Pediatr* 2010;17:1448-50. [\[CrossRef\]](#)
22. Sachwitz D, Hass HJ, Krause H. Bilateral intrauterine testicular torsion in a newborn. *Urologe A* 2012;51:60-2. [\[CrossRef\]](#)
23. Kawamura M, Kuribayashi S, Yamamichi G, Nakano K, Kishimoto N, Tsutahara K, et al. A Case of Prenatal Testicular Torsion. *Hinyokika Kyo* 2016;62:389-91.